

Antiphospholipid Syndrome

What is Antiphospholipid Syndrome (APS)?

APS is a blood disorder caused by abnormal proteins (antibodies) in the blood that leads to an increase risk of blood clotting and problems with pregnancy. APS is diagnosed in people who have blood clots or problems with pregnancy AND who have evidence of these abnormal antibodies in the blood.

What are antibodies?

Antibodies are proteins that are normally produced by the immune system to fight infections. The immune system produces different antibodies for different infections. APS occurs when antibodies are produced by the immune system that are directed against the body rather than infections; it is therefore called an auto(self)immune disorder.

How do you measure these antibodies?

Antibodies in APS are measured using a blood test. There are three blood tests we use;

1. Lupus Anticoagulant (LAC)
2. Anticardiolipin Antibody (ACLA)
3. beta2microglobulin (B2M).

It is important to remember that not everybody who has these antibodies in the blood will develop blood clots; some people have these antibodies but never develop any problems with blood clots. Sometimes these antibodies occur after an infection and then disappear.


How did I get APS?

We are not sure why APS develops. It is more common in females than males.

What are the main problems with having APS?

The main problem with having APS is being at risk of having a blood clot. Blood clots can occur anywhere in the body; often the veins in the leg, sometimes in the arteries of the body and sometimes APS can even cause strokes.

What do I need to do?



If you have APS it is likely you will need blood thinning medication (anticoagulation) to reduce the risk of having a blood clot. Anticoagulation works by stopping the blood clotting system working properly and prevents blood clots in patients with APS.

How long will I have to be on blood thinning for?

Unfortunately, you are at a high risk of having another blood clot if you have already had one blood clot and have blood tests showing you have APS. Blood thinning will reduce this risk significantly – current recommendations suggest that you should stay on blood thinning as long as the risk of blood thinning (bleeding) do not outweigh the benefits. You should talk about continuing blood thinning with your doctor.

What about problems with pregnancy and APS? Is this managed any differently?

APS is definitely associated with problems getting pregnant, early and recurrent miscarriage and pre-term labour. There is some evidence to suggest that treatment with blood thinning (specifically heparin) AND the use of low dose aspirin may reduce the risk of having problems with pregnancy and APS. It may be that aspirin should be started before getting becoming pregnant and heparin should be used once the pregnancy has been confirmed. You will need to talk to your doctor and your obstetrician regarding this treatment.

Resources used to produce this information sheet.

- Rand, H The Antiphospholipid Syndrome. Haematology 2007 pp 136 – 142
- Garda, DA Khamashata, MA Crowther MA. How we diagnose and treat thrombotic manifestations of the Antiphospholipid syndrome; a case based review. Blood 2007 pp 3122 – 3127